

Piedmont Cancer Registry
City of Turin (899.806 residents at the 2001 census)

PROSTATE - ICD-10: C 61

COMMENTS:
Incidence has dramatically increased, due to the spread of PSA practice. Mortality seems not to have been importantly influenced yet. Indeed, survival has increased a lot, but this is in large part due to diagnosis anticipation. Long term survival still decreases, but this is probably due to old age at incidence and to the insufficient statistical adjustment.

INCIDENCE Last Update: 2003-2005

		MEN
N° (average/annual)		723
RATES	Crude	167.3
	Age adj (ITA81)	101.3
	Age adj (WORLD)	69.3

MORTALITY Last Update: 2003-2005

		MEN
N° (average/annual)		132
RATES	Crude	30.6
	Age adj (ITA81)	15.7
	Age adj (WORLD)	10.3

SURVIVAL Last Update: 1995-1999 (f.u. 2004)

		MEN
N° of cases		1963
1 year observed		89.4
1 year relative		93.9
5 year observed		63.3
5 year relative		82.7
5 year standardized		79.0

SURVIVAL Complete data: 1985-1999 (f.u. 2004)

		MEN
N° of cases		5092
10 year observed		34.9
10 year relative		66.2
15 year observed		18.9
15 year relative		54.5

5 year Relative Survival TRENDS (f.u. 2004)

		MEN
1985-1987		52.2
1988-1990		59.8
1991-1993		65.7
1994-1996		76.5
1997-1999		86.6

PREVALENCE 31/12/2004

N° of cases x 100.000

		MEN
observed (since 1985)		969.0
overall (estimated)		1027.6

INDICATORS / METHODS

INCIDENCE - MORTALITY - NUMBER OF CASES - AGE SPECIFIC RATES - CRUDE RATES - STANDARDIZED RATES - SURVIVAL - OBSERVED SURVIVAL - RELATIVE SURVIVAL AND AGE-STANDARDIZED RELATIVE SURVIVAL - PREVALENCE - NOTES

