

Piedmont Cancer Registry
City of Turin (910,504 inhabitants at 31.12.2009)

BLADDER - ICD-10: C 67

COMMENTS:

Incidence is slightly increasing in both sexes. Mortality slowly goes down in men, and it is stable in women. Trends in survival are in part an artefact, due to the change in malignancy definition, as the parallel decrease in mortality in men clearly shows.

INCIDENCE Last Update: 2004-2006

		MEN	WOMEN
N° (average/annual)		413	107
RATES	Crude	95.7	22.7
	Age adj (ITA81)	58.2	13.7
	Age adj (WORLD)	40.3	7.6
Age specific rates		► MEN	► WOMEN

MORTALITY Last Update: 2004-2006

		MEN	WOMEN
N° (average/annual)		87	27
RATES	Crude	20.2	5.7
	Age adj (ITA81)	10.6	2.9
	Age adj (WORLD)	6.9	1.3
Age specific rates		► MEN	► WOMEN

SURVIVAL Last Update: 1995-1999 (f.u. 2004)

		MEN	WOMEN
N° of cases		949	252
1 year observed		79.8	74.6
1 year relative		83.5	77.5
5 year observed		52.7	45.3
5 year relative		67.0	56.3
5 year standardized		67.0	61.0

SURVIVAL Complete data: 1985-1999 (f.u. 2004)

		MEN	WOMEN
N° of cases		3619	928
10 year observed		38.7	36.9
10 year relative		62.4	58.8
15 year observed		26.9	26.2
15 year relative		57.9	55.4

5 year Relative Survival TRENDS (f.u. 2004)

	MEN	WOMEN
1985-1987	72.0	58.4
1988-1990	70.8	74.1
1991-1993	70.9	72.8
1994-1996	67.6	56.9
1997-1999	66.8	56.0

PREVALENCE 31/12/2004

N° of cases x 100.000

	MEN	WOMEN
observed (since 1985)	386.3	84.4
overall (estimated)	456.8	100.6

INDICATORS / METHODS

INCIDENCE - MORTALITY - NUMBER OF CASES
AGE SPECIFIC RATES - CRUDE RATES - STANDARDIZED RATES -
SURVIVAL - OBSERVED SURVIVAL
RELATIVE SURVIVAL AND AGE-STANDARDIZED RELATIVE SURVIVAL -
PREVALENCE - NOTES

